



Contribution ID: 58

Type: Oral

## Evidence for a toroidal electric dipole mode in nuclei

*Friday, 21 July 2023 13:35 (15 minutes)*

A multipole expansion of electromagnetic fields shows that besides the well-known electric and magnetic terms a toroidal dipole mode must exist, where the electric current is bent on the surface of a torus (1). Such modes appear in a wide variety of fields ranging from solid-state physics, quantum and nonlinear optics, to particle physics and astronomy. A toroidal electric mode should also appear in all nuclei as response to an external dipole field, like the isovector giant dipole resonance or the isoscalar compression resonance due to nuclear density variations. Such a mode was predicted more than 50 years ago, but clear experimental evidence is lacking so far (2). Using a combination of high-resolution inelastic scattering experiments with photons (3), electrons (4) and protons (5), we identify for the first time candidates for toroidal excitations in the nucleus  $^{58}\text{Ni}$  and demonstrate that transverse electron scattering form factors represent an unique experimental observable to prove their nature (6). While the present case refers to a nucleus with almost equal proton and neutron number, toroidal excitations might also offer an explanation for the observation of the pygmy dipole resonance at low energies in nuclei with neutron excess (7).

- (1) V.M. Dubovik, V.V. Tugeshev, Phys. Rep. 187, 145 (1990).
- (2) A. Repko et al., Eur. Phys. J. 55, 242 (2019).
- (3) M. Scheck et al., Phys. Rev. C 88, 044304 (2013); J. Sinclair et al., to be published.
- (4) W. Mettner et al., Nucl. Phys. A 473, 160 (1987).
- (5) P. von Neumann-Cosel, A. Tamii, Eur. Phys. J. A 55, 110 (2019).
- (6) V.O. Nesterenko et al., Phys. Rev. Lett. 120, 182501 (2018); Phys. Rev. C 100, 064302 (2019).
- (7) E.G. Lanza et al., Prog. Part. Nucl. Phys. 129, 104006 (2023).

Supported by DFG under contract SFB 1245 (Project ID No. 79384907)

**Primary authors:** Prof. VON NEUMANN-COSEL, P. (Institut für Kernphysik, Technische Universität Darmstadt, Germany); Dr NESTERENKO, V. (Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, Dubna, Russia); Mrs BRANDHERM, I. (Institut für Kernphysik, Technische Universität Darmstadt); Mr VISHNEVSKIY, P.I. (Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, Dubna, Russia); Prof. MATSUBARA, H. (Research Center for Nuclear Physics, Osaka University, Japan); Prof. RICHTER, A. (Institut für Kernphysik, Technische Universität Darmstadt, Germany); Prof. SCHECK, M. (School of Computing, Engineering, and Physical Sciences, University of the West of Scotland, Paisley, UK); Mrs SINCLAIR, J. (School of Computing, Engineering, and Physical Sciences, University of the West of Scotland, Paisley, UK); Prof. TAMII, A. (Research Center for Nuclear Physics, Osaka University, Japan)

**Presenter:** Prof. VON NEUMANN-COSEL, P. (Institut für Kernphysik, Technische Universität Darmstadt, Germany)

**Session Classification:** Session 16

**Track Classification:** Fundamental Symmetries and Interactions