



Contribution ID: 97

Type: **Invited Oral**

## Search for low-lying octupole-isovector excitations

*Monday, 17 July 2023 11:10 (25 minutes)*

Two-component quantum systems develop fundamental modes in terms of in-phase and out-of phase motion of the subsystems. For the nuclear many-body system the framework of the proton-neutron Interacting Boson Approximation (IBA-2) predicted the latter type of excitations as so-called mixed symmetry states. The experimental fingerprint is a strong M1 transition to the proton-neutron symmetric coupled state, which is usually the lowest-lying state for a given spin and parity combination, and a weakly collective decay to the ground state. In this contribution results from  $^{95}\text{Mo}(n,\gamma)$  and  $^{143}\text{Nd}(n,\gamma)$  experiments employing the EX-ILL setup are reported. These measurements aimed to confirm candidates for low-lying octupole isovector (mixed-symmetry) states by measuring the relative intensity and multipole-mixing ratio in the crucial decay to the first  $3^-$  level. Furthermore, for  $^{144}\text{Nd}$  the lifetime of the candidate was remeasured using GAMS. In the talk the resulting experimental picture will be presented and for  $^{96}\text{Mo}$  an alternative interpretation for the observed M1 strength given.

**Primary author:** Dr SCHECK, Marcus (University of the West of Scotland)

**Presenter:** Dr SCHECK, Marcus (University of the West of Scotland)

**Session Classification:** Session 2

**Track Classification:** Experimental Nuclear Structure