



Contribution ID: 164

Type: Oral

Investigation of excited states in ^{76}As of interest for $0\nu\beta\beta$ decay

Thursday, July 20, 2023 6:00 PM (15 minutes)

Since many years the quest for experimental evidences for the neutrino less double beta decay is open. Such measurement would imply the neutrino being identical to its antiparticle, therefore being a Majorana particle. A measurement of the lifetime for this process would lead to a precise estimation of the neutrino mass. However, very precise nuclear physics data are needed to achieve an accurate theoretical prediction. Indeed, the predicted lifetimes and decay rates may vary by many factors depending on the nuclear interaction

chosen, and more and more precise nuclear physics information is needed to match the experimental sensitivity

of planned neutrino experiments. Large experimental efforts have been performed to study, theoretically [1] and experimentally [2, 3], the double beta decay of ^{76}Ge to ^{76}Se . While a crucial role is played by the wavefunctions of the mother ground state and the low-lying states of the daughter, theory shows that a non-negligible role can be played by the structure of the intermediate ^{76}As [4]. An experiment is planned with the FIPPS apparatus at ILL, which aims at an unambiguous identification of the $1+$ levels of ^{76}As . Such identification would help theory constrain the matrix elements of the double beta decay. Such levels will be identified using γ -ray spectroscopy techniques after (n,γ) reactions on ^{75}As . The spin and parity assignment of the excited levels will be possible thanks to angular correlation measurements.

Preliminary results from the FIPPS experiments will be shown as well as the characterization of the performance of the instrument.

[1] J. Menéndez et al.: Commissioning of the Active TARget and Time Projection Chamber Occupancies of individual orbits, and the nuclear matrix element of the ^{76}Ge neutrinoless $\beta\beta$ decay, 2009, Phys. Rev. C.

[2] M. Agostini et al.: Results on neutrinoless double- β decay of ^{76}Ge from phase 1 of the gerda experiment, 2013, Phys. Rev. Lett.

[3] F. ˇSimkovic et al.: Neutrinoless double beta decay of ^{76}Ge , ^{82}Se , ^{100}Mo , and ^{136}Xe to excited $0+$ states, 2001, Phys. Rev. C.

[4] J. T. Suhonen: Value of the axial-vector coupling strength in β and $\beta\beta$ decays: A review., 2017, Frontiers in Physics.

Presenter: DOMENICHETTI1, Lorenzo

Session Classification: Session 13A

Track Classification: Experimental Nuclear Structure