

Metamagnetism & superconductivity in UTe₂

William Knafo¹, Daniel Braithwaite², Stéphane Raymond²,
Michal Valiska^{2,3}, Tristan Thebault¹, Marc Nardone¹,
Abdelaziz Zitouni¹, Alexandre Pourret², Paul Steffens⁴,
Koji Kaneko⁵, Eric Ressouche², Adrien Rosuel²,
Jean-Pascal Brison², Jacques Flouquet², Dai Aoki⁶,
Georg Knebel² and Gérard Lapertot²

¹LNCMI - Laboratoire National des Champs Magnétiques Intenses
143 Avenue de Rangueil, 31400 Toulouse, France

²CEA - Commissariat à l'énergie atomique et aux énergies alternatives
38054 Grenoble Cedex, France

³Charles University, Faculty of Mathematics and Physics
Ke Karlovu 3, 121 16 Praha 2, Czech Republic

⁴ILL - Institut Laue-Langevin
38042 Grenoble Cedex 9, France

⁵JAEA - Japan Atomic Energy Agency
Tokai, Naka, Ibaraki 319-1195, Japan

⁶IMR - Institut for Materials Research
Tohoku University 2-1-1 Katahira, Aoba-ku, Sendai 980-8577, Japan

In 2019, unconventional superconductivity was observed in the heavy-fermion paramagnet UTe₂ and a spin-triplet nature of the superconducting pairing has been proposed for this compound initially presented as a nearly-ferromagnet[1, 2]. Soon after, multiple superconducting phases were found to develop near magnetic transitions in UTe₂ under intense magnetic fields and high pressures[3, 4, 5, 6, 7, 8].

Here, I will present a selection of results performed within a French-Japanese collaboration on UTe₂. Experiments under combined extreme conditions showed that multiple superconducting phases can be induced by a

magnetic field, sometimes coupled with pressure, in the vicinity of metamagnetic transitions [4, 6, 8] (see also [5, 7]). From inelastic neutron scattering at zero magnetic field, we evidenced the presence of quasi-two-dimensional antiferromagnetic fluctuations in UTe₂, which is also a two-legs magnetic ladder [9] (see also [10, 11]). Their gapping in the zero-field superconducting phase indicates that these fluctuations may play a role in the superconducting mechanism [12, 13].

Neutron scattering is a unique tool to microscopically unravel the role of magnetism, and particularly of the magnetic fluctuations, for the development of unconventional superconductivity in correlated-electrons materials. However, metamagnetism in UTe₂ occurs at fields far beyond what is feasible today for inelastic neutron scattering (fields up to 36 T at ambient pressure, or up to 15-20 T under pressure may be needed). As perspectives, I will discuss how the extension of neutron techniques to higher magnetic fields, possibly coupled with very low temperatures and/or high pressures, will constitute a milestone to understand the interplay between magnetism and superconductivity in materials as UTe₂.

References

- [1] S. Ran, C. Eckberg, Q.P. Ding *et al.*, *Science* **365** (2019) 684; DOI: 10.1126/science.aav8645
- [2] D. Aoki, A. Nakamura, F. Honda *et al.*, *J. Phys. Soc. Japan* **88** (2019) 043702; DOI: 10.7566/JPSJ.88.043702
- [3] D. Braithwaite, M. Vališka, G. Knebel *et al.*, *Communication Physics* **2** (2019) 147; DOI: 10.1038/s42005-019-0248-z
- [4] G. Knebel, W. Knafo, A. Pourret *et al.*, *J. Phys. Soc. Japan* **88** (2019) 063707; DOI: 10.7566/JPSJ.88.063707
- [5] S. Ran, I-L. Liu, Y. Suk Eo *et al.*, *Nature Physics* **15** (2019) 1250; DOI: 10.1038/s41567-019-0670-x
- [6] W. Knafo, M. Nardone, M. Vališka *et al.*, *Communication Physics* **4** (2021) 40; DOI: 10.1038/s42005-021-00545-z
- [7] D. Aoki, F. Honda, G. Knebel *et al.*, *J. Phys. Soc. Japan* **89** (2020) 053705; DOI: 10.7566/JPSJ.89.053705
- [8] M. Vališka, W. Knafo, G. Knebel *et al.*, *Phys. Rev. B* **104** (2021) 214507; DOI: 10.1103/PhysRevB.104.214507

- [9] W. Knafo, G. Knebel, P. Steffens *et al.*, *Phys. Rev. B* **104** (2021) L100409; DOI: 10.1103/PhysRevB.104.L100409
- [10] C. Duan, K. Sasmal, M. B. Maple *et al.*, *Phys. Rev. Lett.* bf 125 (2020) 237003; DOI: 10.1103/PhysRevLett.125.237003
- [11] N. P. Butch, S. Ran, S. R. Saha *et al.*, *npj Quantum Materials* **7** (2022) 39; DOI: 10.1038/s41535-022-00445-7
- [12] C. Duan, R. E. Baumbach, A. Podlesnyak *et al.*, *Nature* **600** (2021) 636; DOI: 10.1038/s41586-021-04151-5
- [13] S. Raymond, W. Knafo, G. Knebel *et al.*, *J. Phys. Soc. Japan* **90** (2021) 113706; DOI: 10.7566/JPSJ.90.113706