



Contribution ID: 29

Type: not specified

Measurement of prompt γ -rays from the neutron capture reaction of Pd isotopes in the eV energy range

Thursday, 24 March 2022 09:15 (30 minutes)

In the compound process of neutron-induced nuclear reactions, the parity symmetry (P-violation) is violated due to the effect of the weak interaction. It has been experimentally found that the helicity dependence of the reaction cross section due to the P-violating nucleon-nucleon interaction is enhanced by up to six orders of magnitude compared to the bare effect observed in few-nucleon reactions[1]. This amplification effect has been explained by using a statistical model based on the Random Matrix Theory. In this model, transition matrix element is expected to be inversely proportional to the square-root of the level density N . The accuracy of the existing experimental data of W , however, are not sufficient to verify the model, and thus more accurate data are demanded. W is related to the asymmetry A_L of the helicity-dependent reaction cross section as the following equation;

$$A_L \approx -\frac{2xW}{E_p - E_s} \sqrt{\frac{\Gamma_s^n}{\Gamma_p^n}} \left(x \equiv \sqrt{\frac{\Gamma_{p,j=1/2}^n}{\Gamma_p^n}} \right), \quad (1)$$

where E_s and E_p are the resonance energies of the s-wave and the p-wave resonances, respectively. Γ_s^n and Γ_p^n are the corresponding neutron widths, respectively. x is the ratio of the partial p-wave neutron width to the total neutron width, it can be determined by measuring the angular dependence of the emitted γ -rays of the (n, γ) reaction [2]. Therefore, by measuring A_L and x , one can determine W experimentally from Eq. (1).

In this study, we focus on Pd isotopes which have relatively small values of N , and consequently N dependences of W are rather significant. To obtain x , the angular distributions of the prompt γ -rays from the p-wave resonance of the Pd isotopes were measured at the J-PARC MLF BL04 in February 2021. As a preliminary result, the following values of x were obtained for ^{108}Pd ;

$$x = 0.9986_{-0.0099}^{+0.0003} \quad \text{or} \quad x = -0.9986_{-0.0003}^{+0.0099}. \quad (2)$$

In this contribution, the experimental procedure and the result will be presented.

[1] G. E. Mitchell et al., Phys. Rep. 354, 157 (2001).

[2] V. V. Flambaum et al., Nucl. Phys. A 435, 352 (1985).

Primary authors: Mr YOSHIKAWA, Hiromoto (Osaka University); Mr ABE, Ryota (Nagoya University); Mr ISHIZAKI, Kohei (Nagoya University); Mr ITO, Yuki (Nagoya University); Mr ENDO, Shunsuke (JAEA); Dr OKUDAIRA, Takuya (Nagoya University); Mr KAMEDA, Kento (Tokyo Institute of Technology); Dr KITAGUCHI, Masaaki (Nagoya University); Dr KIMURA, Atsushi (JAEA); Dr SAKAI, Kenji (JAEA); Dr SHIMA, Tatsushi (Osaka University); Prof. SHIMIZU, Hirohiko (Nagoya University); Mr TAKADA, Shusuke (Kyushu University); Mr NAKAI, Asuka (Osaka University); Dr HIROTA, Katsuya (KEK); Mr FUJIIIE, Takuhiro (Nagoya University); Dr FUJIOKA, Hiroyuki (Tokyo Institute of Technology); Dr YOSHIOKA, Tamaki (Kyushu University)

Presenter: Mr YOSHIKAWA, Hiromoto (Osaka University)