

Contribution ID: 23

Type: not specified

Lifetime measurements in the sub-picosecond regime with the py-coincidence Doppler-shift attenuation method

Thursday, 24 March 2022 11:00 (30 minutes)

Nuclear level lifetimes are important properties of the atomic nucleus, as they yield information about transition strengths and nuclear wave functions. An established method to determine lifetimes in the sub-picosecond regime is the particle- γ coincidence Doppler-shift attenuation method (p γ -DSAM) [1]. As opposed to most DSAM approaches, the additional coincident detection of emitted γ -ray and scattered projectile allows the selection of excited states, thus eliminating feeding contributions. The coincidence data is obtained with the SONIC@HORUS detector array [2] situated at the 10 MV FN tandem accelerator in Cologne. It consists of 12 silicon particle detectors and 14 high-purity germanium (HPGe) detectors for the detection of γ -rays. With this method and setup, lifetimes of several dozens of nuclear levels can be determined in a single experiment. Lately, the DSA method has been used to determine lifetimes of excited low-spin states of nuclei in the mass region of A \approx 100 and above. Recent results on Ru, Sn [3] and Te [4] isotopes will be presented in this contribution.

Supported by the DFG (ZI-510/9-1).

- [1] A. Hennig et al., NIM A 794, 171 (2015).
- [2] S.G. Pickstone et al., NIM A 875, 104 (2017).
- [3] M. Spieker et al., Phys. Rev. C 97, 054319 (2018).
- [4] S. Prill et al., accepted at Phys. Rev. C

Primary author: Ms PRILL, Sarah (University of Cologne, Institute for Nuclear Physics, Cologne, Germany)

Co-authors: Ms BOHN, Anna (University of Cologne, Institute for Nuclear Physics, Germany); Ms DEKE, Christina (University of Cologne, Institute for Nuclear Physics, Cologne, Germany); Mr HEIM, Felix (University of Cologne, Institute for Nuclear Physics, Cologne, Germany); Mr WEINERT, Michael (University of Cologne, Institute for Nuclear Physics, Cologne, Germany); Prof. ZILGES, Andreas (University of Cologne, Institute for Nuclear Physics, Cologne, Germany); Mr WEINERT, Michael (University of Cologne, Institute for Nuclear Physics, Cologne, Germany); Prof. ZILGES, Andreas (University of Cologne, Institute for Nuclear Physics, Cologne, Germany); Prof. ZILGES, Andreas (University of Cologne, Institute for Nuclear Physics, Cologne, Germany); Prof. ZILGES, Andreas (University of Cologne, Institute for Nuclear Physics, Cologne, Germany); Prof. ZILGES, Andreas (University of Cologne, Institute for Nuclear Physics, Cologne, Germany); Prof. ZILGES, Andreas (University of Cologne, Institute for Nuclear Physics, Cologne, Germany); Prof. ZILGES, Andreas (University of Cologne, Institute for Nuclear Physics, Cologne, Germany); Prof. ZILGES, Andreas (University of Cologne, Institute for Nuclear Physics, Cologne, Germany); Prof. ZILGES, Physics, Cologne, Institute for Nuclear Physics, Cologne, Germany); Prof. ZILGES, Physics, Cologne, Institute for Nuclear Physics, Cologne, Institute for Nuclear

Presenter: Ms PRILL, Sarah (University of Cologne, Institute for Nuclear Physics, Cologne, Germany)