

# Soft Matter Studies with X-rays

Theyengeri Narayanan

ESRF – The European Synchrotron

M. Mitov, *Sensitive Matter - Foams, Gels, Liquid Crystals, and Other Miracles* (Harvard University Press, 2012)

R. Piazza, *Soft Matter: The Stuff that Dreams are Made of* (Springer, 2011)

M. Doi, *Soft Matter Physics* (OUP Oxford, 2013)

R.A.L. Jones, *Soft Condensed Matter* (OUP Oxford, 2004)

LS Hirst, *Fundamentals of soft matter science* (CRC press, 2020)

W. de Jeu, *Basic X-ray Scattering for Soft Matter* (OUP Oxford 2016)

T. Narayanan and O. Konovalov, *Materials*, **13**, 752 (2020);

<https://doi.org/10.3390/ma13030752>

# Outline

- What is Soft Matter?
- Some general features
- Different X-ray techniques employed
- Self-assembly & complexity
- Out-of-equilibrium phenomena
- Summary and outlook

# What is Soft Matter?

**Soft matter** is a subfield of condensed matter physics (CMP) comprising a variety of physical states that are easily deformed by thermal stresses or thermal fluctuations. They include liquids, colloids, polymers, foams, gels, granular materials, and a number of biological materials. These materials share an important common feature in that predominant physical behaviors occur at an energy scale comparable with room temperature thermal energy. At these temperatures, quantum aspects are generally unimportant. Pierre-Gilles de Gennes, who has been called the "founding father of soft matter," received the Nobel Prize in physics in 1991 for discovering that the order parameter from simple thermodynamic systems can be applied to the more complex cases found in soft matter, in particular, to the behaviors of liquid crystals and polymers.

*Matière molle* » Madeleine Veyssié


Today soft matter science is an interdisciplinary field of research where traditional borders between physics and its neighboring sciences such as chemistry, biology, chemical engineering and materials science have disappeared. It is one of the frontiers of CMP along with strongly correlated electron systems and nanoscience.

# Soft Matter: Encounter in everyday life



Sustainable development through more rational design of consumer products

# Soft Matter Systems



Meso-scale structures determine material properties

**SAXS, WAXS, USAXS, GISAXS  
(SANS, USANS, GISANS, etc.)**

# Soft Matter Features

Materials which are soft to touch – characterized by a small elastic modulus (energy/characteristic volume), typically  $10^9 - 10^{12}$  times lower than an atomic solid like aluminum.

Dominance of entropy

Strong influence of thermal fluctuations ( $\sim k_B T$ )

Characteristic size scale or microstructure  $\sim 100 - 1000$  nm

Shear modulus,  $G \sim \text{Energy/Free volume} \gg 10^9 - 10^{12}$  smaller

Low shear modulus ( $G$ )  $\gg$  soft and viscoelastic

Soft Matter studies seek to address the link between microscopic structure/interactions/dynamics and macroscopic properties.

Soft Matter constitutes a significant fraction of modern day Nanoscience/Nanotechnology.

# Self-Assembled Soft Matter Systems


How are these complexes formed?

Kinetic pathways

Biomolecules

How can these complexes be tuned and manipulated?

Functional materials




T. Narayanan et al., Crystallogr. Rev. (2017)


# Soft Matter: COVID-19

COVID-19 pandemic exposed the knowledge-gap

Interaction of detergents with SARS CoV-2



W.C.K. Poon *et al.*, *Soft Matter* (2020)



Formulation of efficient detergents, vaccines, drugs, etc.

# Synchrotron Techniques used in Soft Matter

# Synchrotron Radiation Studies of Soft Matter

- **High spectral brilliance or brightness**

Real time studies in the millisecond range, micro/nano focusing and high  $q$  resolution

Time-resolved SAXS, WAXS, micro-SAXS, USAXS, etc.

- **Partial coherence**

Equilibrium dynamics using the coherent photon flux (for concentrated systems)

Photon correlation spectroscopy (XPCS)

- **Continuous variation of incident energy**

Contrast variation of certain heavier elements, e.g. Fe, Cu, Se, Br, Rb, Sr, etc.

Anomalous Scattering – contrast variation



- **Complementary imaging techniques**

X-ray microscopy, micro and nano tomography, etc.

**Radiation damage**

# Small-Angle X-ray Scattering (SAXS)

Scattering originates from the spatial fluctuations of electron density



$$q = \frac{4\pi}{\lambda} \sin(\theta / 2)$$

Measured Intensity:  $I_S = i_0 T_r \varepsilon \Delta\Omega \left( \frac{d\sigma}{d\Omega} \right)$  Differential scattering cross-section


$i_0$  - incident flux  
 $T_r$  - transmission  
 $\varepsilon$  - efficiency  
 $\Delta\Omega$  - solid angle

$$I(q) = \frac{d\Sigma}{d\Omega} = \frac{1}{V_{Scat}} \frac{d\sigma}{d\Omega}$$

# Ultra SAXS/SAXS/WAXS


Beamline ID02

34 m





$2\pi/q$  (nm)




Time resolution: < 100 μs



# Size scales probed by SAXS & related techniques



# SAXS from dilute spherical particles






Modeling or simulation required to extract quantitative information

# Protein Solution Scattering

Traditionally a few structural features: size, shape, size or density distribution

Beamline - BM29




# Form & Structure Factors

Differential scattering cross-section per unit volume



$$I(q) = N(\Delta\rho^* V)^2 P(q) S_M(q)$$

FT of radial electron density






pair correlation function

$S(q)$  from liquid state theories (e.g. Percus-Yevick (PY) ) or simulations




# Structure Factors at high packing fractions

E.g. 60%




Glass




Crystal

# High Resolution USAXS




# SAXS by a Partially Coherent Beam


Standard beam and area detector



Highly collimated beam and high resolution detector



$$\text{Speckle size} \sim \frac{\lambda d_{SD}}{\sigma}$$



$$\xi_T \sim \frac{\lambda R_S}{d} \sim (10 - 100 \mu\text{m})$$

$$\xi_L = \frac{c}{\Delta\nu} = \lambda \left( \frac{\lambda}{\Delta\lambda} \right) \sim 1 \mu\text{m}$$



# X-ray Photon Correlation Spectroscopy (XPCS)

XPCS at small and wide angles: **ID10 beamline** (Y. Chushkin)

At ultra low angles,  $10^{-3} \leq q \leq 10^{-1} \text{ nm}^{-1}$  : ID02 beamline

Analogous to dynamic light scattering

Dilute silica colloids of 600 nm in size



$$\langle \Delta r^2(\tau) \rangle = 6D_0\tau$$

mean-square displacement

$$D_0 = \frac{k_B T}{6\pi\eta R}$$


diffusion constant  
(Stokes-Einstein)

**Multi-speckle XPCS**

Suitable for optically opaque systems

# Multi-Speckle XPCS


Dilute silica colloids of 600 nm in size



Ensemble averaged  
Intensity autocorrelation function

$$g_2(\mathbf{q}, t) = \frac{\langle I(\mathbf{q}, \tau) I(\mathbf{q}, \tau + t) \rangle}{\langle I(\mathbf{q}) \rangle^2} = 1 + \beta g_1(\mathbf{q}, t)^2$$

Speckle contrast:  $\beta > 0.4$



Brownian dynamics:  $\Gamma(q) = D_0 q^2$


Direction dependent analysis

# Grazing Incidence Small-Angle X-ray Scattering

ID10 Beamline (O. Kononov)

(GISAXS)

$$q_{x,y,z} = \frac{2\pi}{\lambda} \begin{cases} \cos \alpha_f \cos \theta_f - \cos \alpha_i \cos \theta_i \\ \cos \alpha_f \sin \theta_f - \cos \alpha_i \sin \theta_i \\ \sin \alpha_i + \sin \alpha_f \end{cases}$$



# ID10 Surface & Interface Scattering Beamline

GISAXS, GID, XRR, GIXF

(O. Konovalov)


Multipurpose instrument for surface/interface studies

4 circle diffractometer

Beam deflector stage for liquid surfaces

The two-crystal  
deflector stage rotates  
the X-ray beam around  
a fixed point on the  
liquid surface



$\left\{ \begin{array}{l} q_x \\ q_y \\ q_z \end{array} \right.$




# Soft Interfaces Scattering

Beamline ID10


Using higher energy X-rays  
( $> 30$  keV)




Varying the penetration depth




# SAXS/WAXS from Semi-crystalline Polymers




Spherulite morphology



# Scanning Micro-diffraction on HDPE spherulites



- **high density poly-ethylene**
- spherulites under polarized light banded structures indicating long range order






- **SAXS/WAXS** patterns
- line scans across the center reveal information on crystallite orientation

*M. Rosenthal et al., Angewandte Chemie, 123, 9043-9047 (2011)*

# Chirality of twisted polymer crystals

Azimuth/Intensity vs Distance from the center in  $\mu\text{m}$



- $35^\circ$  tilt between c-axis and the normal of the base plane of crystalline lamellas
- orientation of b-axis aligned with growth direction
- chirality can be determined

# Soft Matter Self-Assembly


# Spontaneous self-assembly of micelles and vesicles

E.g. surfactants, lipids or block copolymers

Large variety of equilibrium structures

Dynamics of formation is very little explored

## Self-assembly of micelles and vesicles



Rate-limiting steps » predictive capability


Kinetic pathway: stopped-flow rapid mixing & time-resolved SAXS

# Triggering & Synchronization of Dynamic Processes

E.g. concentration/pH jump (rapid mixing)



Rapid temperature or pressure change

Flash photolysis



# Micelle – Vesicle Transformation

## Spontaneous self-assembly of small unilamellar vesicles



At the closing state:  $R_{\max} \approx \frac{4(2\kappa + \bar{\kappa})}{\Lambda}$




$\kappa$  &  $\bar{\kappa}$  - bending moduli  
 $\Lambda$  - line tension

The whole evolution of the scattering curves can be described by a mechanistic model

# Hepatitis B Virus Capsid Assembly

Prof. U. Raviv, HUJI

Umbrella sampling of MC Simulation:





A comprehensive library of distinct intermediates


Configurational entropy favors holey capsid structures

# Pathways of Hepatitis B Virus Capsid Assembly

Simultaneous change of ionic strength and temperature





Physiological conditions




R. Asor *et al.*, JACS (2020)

# Pathways of Hepatitis B Virus Capsid Assembly


Aggressive conditions



Mild conditions




Aggressive conditions




Dimer-dimer association free energy controls the earliest steps of the reaction, which dictates the subsequent assembly pathway

# Multi-step hierarchical self-assembly of microtubules



Simple ingredients, a prototypical surfactant (SDS) and a naturally abundant polysaccharide ( $\beta$ -cyclodextrin) in water forming complex hierarchical structures.

$2\beta$ -CD+SDS @ 75 °C  $\rightarrow$  25 °C




L. Jiang et al., *Soft Matter* (2011)

Spectacular self-assembly spanning size scales of 3 orders leading to formation of microtubules with a diameter of about 1.2  $\mu$ m


J. Landman *et al.*, *Science Advances* (2011)

# Nucleation and Growth Mechanism



Central radius & wall thickness



Concentration scaling



Inward growth of tubes by nucleation & growth




# Self-Assembly of 2D Superlattices

Formation mechanism of two-dimensional superlattices from PbSe nanocrystals at vapour/liquid interface

WAXS: Orientation/domain size

J.J. Geuchies, *et al.*, Nature Materials (2016)

SAXS: Lateral organization



Hexagonal array

Deformed array

Square lattice

Crystalline bridges between the nanocrystals


# Out-of-equilibrium dynamics

# Soft Matter: Out-of-equilibrium Dynamics

Out-of-equilibrium dynamics of systems far away from equilibrium


## Multi-speckle XPCS

Series of scattering patterns




$$c_I(t, \tau) = \frac{\langle I_p(t) I_p(t + \tau) \rangle_p}{\langle I_p(t) \rangle_p \langle I_p(t + \tau) \rangle_p}$$

Time resolved correlation function



$$g^{(2)}(Q, \tau) = \langle c_I(t, \tau) \rangle_t$$





# Dynamics of Active Particles

Active matter: understanding the physics of life from complex systems perspective

## Complex systems



## Self-propelled particles





Micro-swimmers (microorganisms or Janus particles in a catalytic medium)

## Diffusiophoresis


Probe the emergent dynamics by XPCS

Control parameter, Péclet number ( $Pe$ )

# Multispeckle XPCS Study of Active Dynamics



## Self-propulsion & associated dynamics



Weaker  $q$ -dependence – tendency towards superdiffusive behavior

# Emergent Active Dynamics

propulsive

diffusive

transit

Gaussian

$$(\Delta v \cdot q)^2$$

Exponential



$$Dq^2$$

Gaussian

$$v$$

$$|g_1(q, t)|^2 = |g_{1,A}(q, t)|^2 \times |g_{1,D}(q, t)|^2 \times |g_{1,T}(t)|^2$$

Mean velocity ( $v$ ) and its variance ( $\delta v$ )  
Effective diffusion coefficient ( $D_{\text{eff}}$ )



Large  $\delta v$  – strong number fluctuations  
Weaker  $q$ -dependence of ( $\Gamma_e$ )

# Summary & Outlook

- High brilliance X-ray scattering is a powerful method to elucidate the non-equilibrium structure & dynamics of soft matter.
- Time-resolved scattering experiments in the sub-millisecond range can be performed even with dilute samples.
- **Combination of nanoscale spatial and millisecond time resolution makes synchrotron techniques unique in these studies.**
- Experiments can be performed in the functional state of the system.
- Challenges lie in the ability to investigate multicomponent systems and radiation sensitive specimen.
- The emphasis has become on quantitative studies of highly complex systems by exploiting the coherence properties of extremely bright synchrotron sources. In particular to problems related to the Physics of Life.