



Contribution ID: 85

Type: Oral

## Simultaneous Interfacial Rheology and Neutron Reflectometry studies on fluid interfaces

*Tuesday, 16 July 2024 16:50 (20 minutes)*

Fluid interfaces with adsorbed substances are present in many systems in nature and industrial processes. Such interfaces often have a complex structural configuration which confers them the capability to withstand deformations. Consequently, they have been the object of study in recent decades both from a structural and rheological point of view. To date, there have been limited examples of simultaneous measurements of the interfacial rheology and the structure of complex fluid interfaces. This aspect holds particular significance, considering the challenge of comparing independently conducted structural and rheological experiments, where reproducing identical experimental conditions, such as temperature and/or concentration and/or compression history for monolayer at the air-water interface, is difficult. Specifically, there is great interest in the study of Langmuir monolayers of fatty acids and phospholipids which appear in many biophysical processes. This work focuses on the development, building, and exploitation of an interfacial shear rheometer, with DWR geometry, to be used on the neutron horizontal reflectometer FIGARO at Institut Laue-Langevin (ILL). Consequently, the instrument allows for simultaneous measurements of neutron reflectometry and interfacial rheology. In particular, a DWR probe 3D printed in titanium has been commissioned for the Anton Paar MCR702e Space rheometer available at the PSCM, and a suitable shear channel with annular geometry (machined in PTFE) has been designed and built to be used in the Langmuir trough. Notably, an improved Flow Field-Based data analysis software package has been developed to properly subtract bulk phases contribution, taking into account non-linear velocity profiles. The performance of the new instrument is illustrated with a study of the isothermal compression of C19 fatty acid Langmuir monolayers (FALMs). Additionally, Brewster Angle Microscopy (BAM) has been used to observe the formation of structures above the micron scale at the interface. The studies carried out attempt to shed light on the mechanism of loss of molecules observed in condensed phases at high interfacial pressures in monolayers of fatty acids from a dynamic and structural point of view.

### Please select the related topic from the list below

Dynamics of surfaces, interfaces, and nanostructures

**Primary authors:** MAESTRO, Armando (Materials Physics Center); CARRASCOSA TEJEDOR, Javier (Institut Laue-Langevin); Prof. TAJUELO, Javier (UNED); Prof. RODRIGUEZ-HAKIM, Mariana (UNED); Prof. RUBIO ALVAREZ, Miguel Angel (UNED); SANCHEZ PUGA, Pablo (ILL); GUTFREUND, philipp (Institut Laue-Langevin, Grenoble, France)

**Presenter:** SANCHEZ PUGA, Pablo (ILL)

**Session Classification:** Dynamics of surfaces, interfaces, and nanostructures