




Exploring magnetic correlation lengths in frustrated systems via single-crystal neutron diffraction

Oleg Petrenko
University of Warwick
UK

Outline

- Rare-earth disilicates magnets $\text{RE}_2\text{Si}_2\text{O}_7$ (RE = Er, Ho, Tm)
- Honeycomb/zig-zag antiferromagnets SrRE_2O_4 (RE = Er, Gd, Ho, Dy, Nd, Tb)
- Shastry-Sutherland lattice REB_4 tetraborides (RE = Ho, Tm, Er, Nd)



Acknowledgements

$\text{RE}_2\text{Si}_2\text{O}_7$

M. Islam, M. Ciomaga Hatnean (Warwick & PSI)

N. d'Ambrumenil, G. Balakrishnan (Warwick)

D.D. Khalyavin, P. Manuel, F. Orlandi (ISIS)

J. Ollivier, B. Ouladdiaf (ILL)

SrRE_2O_4

O. Young, S.X.M. Riberolles, G. Balakrishnan (Warwick)

P. Manuel, D.D. Khalyavin (ISIS)

N. Qureshi, O. Fabelo, B. Ouladdiaf, A.R. Wildes (ILL)

REB_4

D. Brunt, G. Balakrishnan, D.A. Mayoh, M.R. Lees (Warwick)

D.D. Khalyavin, P. Manuel, F. Orlandi (ISIS)

R.D. Johnson (UCL)

N. Qureshi, B. Ouladdiaf, A.R. Wildes (ILL)




Fig. 1. Polymorphism of the rare-earth disilicates as observed during the synthesis at normal pressure. Seven different structure types, A ∇ , B $\circ$, C*, D ∇ , E $\bullet$, F*, G*, were identified by X-ray work. Arrow on top of each column indicates melting point of the compound.




Fig. 2. Graphical presentation of the unit-cell geometry of the 7 observed structure types, A-G, each represented by data of one disilicate compound from TABLE I. (Type A ∇ , B $\circ$, C*, D ∇ , E $\bullet$, F*, and G*.)

J. Less-Common Metals, 21 (1970) 1-14

Rare-earth disilicates

$RE_2Si_2O_7$

M. Ciomaga Hatnean et al,
Cryst. Growth Des. 2020, 20, 6636–6648

Single crystal sample are available for:

$Er_2Si_2O_7$ D-type
 $P2_1/b$ space group

$Tm_2Si_2O_7$ C-type
 $C2/m$ space group

$Ho_2Si_2O_7$ E-type
 $Pna2_1$ space group

Rare-earth disilicates

4

J. FELSCHE




Fig. 1. Polymorphism of the rare-earth disilicates as observed during the synthesis at normal pressure. Seven different structure types, A ∇ , B $\circ$, C $\star$, D ∇ , E $\bullet$, F $\star$, G $\star$, were identified by X-ray work. Arrow on top of each column indicates melting point of the compound.





Fig. 2. Graphical presentation of the unit-cell geometry of the 7 observed structure types, A–G, each represented by data of one disilicate compound from TABLE I. (Type A ∇ , B $\circ$, C $\star$, D ∇ , E $\bullet$, F $\star$, and G $\star$.)

J. Less-Common Metals, 21 (1970) 1–14

Er₂Si₂O₇





D-type Er₂Si₂O₇



monoclinic space group P2₁/b

$a=4.69$ Å, $b=5.56$ Å, $c=10.79$ Å, $\gamma=96^\circ$



$T_N=1.85$ K



Er₂Si₂O₇





Er₂Si₂O₇



Magnetic structure, $\mathbf{k} = 0$

Ion	Positions (a, b, c) basis			μ_a	μ_b [μ_B]	μ_c	Moment [μ_B]
1	0.891	0.096	0.351	-5.621	2.105	2.082	6.546
2	0.109	0.404	0.851	-5.621	2.105	-2.082	6.546
3	0.109	0.904	0.649	5.621	-2.105	-2.082	6.546
4	0.891	0.596	0.149	5.621	-2.105	2.082	6.546




Er₂Si₂O₇




Magnetic structure, $\mathbf{k} = 0$

Ion	Positions (a, b, c) basis			μ_a	μ_b [μ_B]	μ_c	Moment [μ_B]
1	0.891	0.096	0.351	-5.621	2.105	2.082	6.546
2	0.109	0.404	0.851	-5.621	2.105	-2.082	6.546
3	0.109	0.904	0.649	5.621	-2.105	-2.082	6.546
4	0.891	0.596	0.149	5.621	-2.105	2.082	6.546


Er₂Si₂O₇



Er₂Si₂O₇



M.J.M. Leask *et al.*, J. Phys. C Solid State **19**, 1173 (1986)



Magnetic properties of D-type Er₂Si₂O₇

1177






Figure 3. The magnetic moment of a single crystal of D-type Er₂Si₂O₇ in μ_B/atom as a function of the magnetic field (in T) applied parallel to the (Δ) a axis and ($\bullet$) 'm' axis at 0.5 K.

Er₂Si₂O₇



$T = 0.24$ K
 $H // a$ axis

$\text{Er}_2\text{Si}_2\text{O}_7$



$T = 0.24 \text{ K}$


$H \parallel a$ axis

Er₂Si₂O₇

$$\mathbf{q} = (0 \ 1/2 \ 1/2)$$



$$\mathbf{q} = (0 \ 1/4 \ 1/4)$$

$$\mathbf{q} = (0 \ 1/3 \ 0)$$





$$T = 0.24 \text{ K}$$

$H // a$ axis



Er₂Si₂O₇




D-type Er₂Si₂O₇ is an Ising-like antiferromagnet with 4-sublattice $\mathbf{k} = 0$ structure below $T_N = 1.85$ K with two pairs of collinear antiferromagnetic moments within the unit cell.

For $H//a$, a narrow and T-sensitive magnetisation plateau exists at 1/3 of saturation value.

The magnetic unit cell undergoes a significant increase within the plateau regime demonstrated by the appearance of a scattering signal at non-integer positions described by the propagation vectors $(0 \ 1/2 \ 1/2)$, $(0 \ 1/4 \ 1/4)$, $(0 \ 1/3 \ 0)$.

The intensity, shape, and width of these field-induced features change rapidly with the applied magnetic field within the magnetisation plateau. The diffraction pattern then returns to integer peaks at higher fields, upon its entrance into the paramagnetic phase and out of the plateau.





D-type Er₂Si₂O₇ is an Ising-like antiferromagnet with 4-sublattice $\mathbf{k} = 0$ structure below $T_N = 1.85$ K with two pairs of collinear antiferromagnetic moments within the unit cell.





For $H//a$, a narrow and T-sensitive magnetisation plateau exists at 1/3 of saturation value.

The magnetic unit cell undergoes a significant increase within the plateau regime demonstrated by the appearance of a scattering signal at non-integer positions described by the propagation vectors $(0 \ 1/2 \ 1/2)$, $(0 \ 1/4 \ 1/4)$, $(0 \ 1/3 \ 0)$.

The intensity, shape, and width of these field-induced features change rapidly with the applied magnetic field within the magnetisation plateau. The diffraction pattern then returns to integer peaks at higher fields, upon its entrance into the paramagnetic phase and out of the plateau.




SrRE₂O₄ (RE= Gd, Ho, Er, Dy, Nd, Tb)




Coexistence of the two different magnetic components:


1/3 magnetisation plateaux for H//a.



Coexistence of the two different magnetic components:
metamagnetic transition for $H//c$.



SrEr_2O_4




For SrEr₂O₄, single-crystal neutron diffraction reveals the nature of field-induced transitions.

For H//a, the applied field induces a transition to an extended, but still not fully long-range ordered UUD state of the Er2 sites.

For H//c, the applied field destroys the long-range antiferromagnetic order of the Er1 sites and induces a much shorter-range order during the metamagnetic transition prior to establishing a long-range ferromagnetic-like state of the Er1 moments.


REB₄ (RE = Ho, Tm, Er, Nd)



Shastry-Sutherland lattice REB_4 tetraborides


REB₄

LINDA YE, TAKEHITO SUZUKI, AND JOSEPH G. CHECKELSKY




ErB₄ and TmB₄

REB₄



TmB₄


REB₄



TmB₄

PRB **102**, 060407(R) (2020)

20





HoB₄



PRB **95**, 024410 (2017)

Scientific Reports 8, 232 (2018)

REB₄




HoB₄




Magnetic susceptibility of NdB₄ in fields applied along and normal to the c axis. The insets show the inverse susceptibility.


$$T_{N1} = 17 \text{ K}, T_{N2} = 6.8 \text{ K}, T_{N3} = 4.9 \text{ K}$$




Field dependent magnetisation curves of NdB₄ for $H \parallel c$.





Magnetisation curves for NdB₄ in a pulsed magnetic field up to 500 kOe.



Magnetic phase diagrams of NdB₄ for different field directions.





LT

$$\mathbf{k}_1 = 0$$

$$\mathbf{k}_2 = (1/5 \ 0 \ -2/5)$$

$$\mathbf{k}_3 = 2\mathbf{k}_2 = (2/5 \ 0 \ -4/5)$$


IT

$$\mathbf{k}_1 = 0$$



$$\mathbf{k}'_2 = (0.142 \ 0.142 \ -2/5)$$

HT



$$\mathbf{k}_1 = 0$$




NdB₄




$$\mathbf{k}_1 = 0$$



NdB₄



In-plane (all-in-all-out)
Fourier component, $m^{k^1}_{xy}$




Out-of-plane Fourier
component, $m^{k^1}_z$,
defining the alternating
nominally up direction of
the internal staggered
field, H_{int}

$$\mathbf{k}_1 = 0$$

$$\mathbf{k}_2 = (1/5 \ 0 \ -2/5)$$

$$\mathbf{k}_3 = 2\mathbf{k}_2 = (2/5 \ 0 \ -4/5)$$



@ 1.5 K

$$m^{k^1}_{xy} = 1.60 \mu_B$$

$$m^{k^1}_z = 0.93 \mu_B$$

$$m^{k^2}_z = 1.72 \mu_B$$





$$m^{k^3}_z = 0.66 \mu_B$$

An anharmonic spin
density wave consisting of
three Fourier components,
 $m^{k^1}_z$, $m^{k^2}_z$, and $m^{k^3}_z$.

NdB₄

$$J_1^x = J_1^y = 0$$





$$J_1^z \neq 0$$




$$\delta^{k1}$$

$$\delta^{k1} m_{xy}^{k1} m_z^{k1}$$



$$\delta^{k1} m_{xy}^{k1} = H_{int}$$



- Ground state of NdB₄ represents a unique interplay between strongly anisotropic spins in a frustrated exchange topology and non-magnetic order parameter
- It involves coexisting instabilities coupled via an internal field created by the non-magnetic distortion, undetectable in the present diffraction data
- One instability is the orthogonal all-in all-out order of the *xy*-components of the moments. Another is SDW of the out-of-plane moments, locked to 5-fold periodicity of the lattice to form 3U2D structure



ErB₄



ErB₄

